

# Kegel Exercises

---

Kegel exercises are designed to strengthen the pelvic floor muscles. These muscles support the bladder and bowel openings in both men and women. Strengthening the muscles of the pelvic floor can aid in preventing leakage of urine or feces with coughing, sneezing, lifting, and other stressful movements. Other benefits of kegels include enhanced sexual function, conditioned muscles to make childbirth easier, decrease and/or prevent prolapse of pelvic organs, and improve the ability to pass stool.

## Who should do Kegel Exercises?

- Women and men with urinary and /or bowel incontinence
- Women and men who have demonstrated weakening of the pelvic floor
- Pregnant women or women who have previously had children
- Middle aged and older women

## What you need to know about Kegel Exercises

The success of Kegel exercises depends on the use of proper technique as well as compliance to a regular exercise program. When doing the exercises, it is important to identify the correct muscles of the pelvic floor. At first, most people contract the abdominal or thigh muscles while neglecting the pelvic floor muscles. This may actually worsen pelvic floor tone and incontinence.

Kegels may be done as a part of biofeedback, when some patients require assistance. Biofeedback consists of placing a sensor on the abdomen and around the anal area, which measure the contraction of the pelvic floor muscles. Occasionally, an electrode may be placed in the vagina in women, or in the rectum in men.

## **Three Steps to an Effective Kegel**

- Learn to tighten the muscles around the vaginal/anal area
- Contract the vaginal and rectal muscles. Note that when you perform steps 1 and 2 correctly, you should also feel the muscles around the anus tighten slightly. This is normal, but do not consciously try to tighten those muscles.
- In a quiet, relaxed setting with no distractions, practice your Kegels and determine how long you can hold your contraction and how many you can do before becoming fatigued. Do not do more than 5-10 reps at time with a 3-5 second hold.

## **Detailed Instructions on how to do Kegels:**

### **Step #1: The urethra—The Functional Stop Test**

After partially emptying your bladder, stop your urine flow in a slow controlled manner, paying attention to how it feels.

**Note:** Stopping and starting the urine repeatedly as an exercise can be harmful and should not be done. Holding your urine can contribute to bladder infection, causing damage to normal urinary reflexes. The Functional Stop Test is useful for assessing what it feels like to gradually stop urine flow, and should not be done more than once per urination.

### **Step #2: The vagina and rectum as your window to the pelvic floor**

For this step, you will internally assess the ability to squeeze and elevate the muscles around the opening of the vagina or rectum, while lying on your back. This is accomplished by inserting one finger into the vaginal or rectal opening while you try to lift and squeeze with the muscles. Pay attention to how this feels.

**Note:** For those who may be unable to check internally, you can place the pad of one finger on the perineal body (the area between the vagina and the anus) and assess for lift in this manner. The internal assessment is preferable, however, because it gives you a better sense of what you are capable of doing with your pelvic floor and provides a way to assess the strength and control you gain as you progress with your Kegels.

### **Step #3: Putting it all together**

Begin practicing while lying on your back with your knees bent, or lying back with pillows under your head and shoulders. When your knees are bent you should have pillows under them so that the muscles around the hips and buttocks can relax. In this position, you can practice the actual Kegel, which involves lifting and tightening the pelvic floor all at one time. This will be a combination of what you felt when you gradually stopped urine flow, with the feeling squeezing and elevating the muscles around the vaginal or rectal area.

### **How long and how many?**

When you are first starting out, it is common to find that you can only hold a Kegel for 3-6 seconds, and that performing 3-6 repetitions causes muscle fatigue. When this happens, you are usually performing the exercise correctly. If you find you can hold it for much longer right away, recheck your technique. For patients with incontinence or pelvic floor weakness, it is important to concentrate on correctly performing the technique and gradually improving the muscle quality and tone, even though you may be weak first starting out.

### **Making Kegels a part of your life**

After you have become experienced in performing these exercises, you will want to do them as follows:

- A series of strong, steady squeezes for 8-10 seconds each

- Try to do 8-10 at a time
- For most people, 30 repetitions per day are adequate.

As they get easier, you will find you can do them while sitting or standing, for example, you can do them while driving, working on the computer, or waiting in long lines.

Please note that you may not see any noticeable change in bowel / bladder control for a while. In some cases it may be many months.

Disclaimer: This document is not intended to take the place of the care and attention of your personal physician or other professional medical services. Talk with your doctor if you have Questions about individual health concerns or specific treatment options.

©2011 The Regents of the University of Michigan  
Author: Bruce M. LaBrecque, RN, PT, U of M PT & MBCP  
Last Revised 06/2011