

A M El-Tawil, MSc, MRCS, PhD, Series Editor

Current management of fecal incontinence: Choosing amongst treatment options to optimize outcomes

Julie Ann M Van Koughnett, Steven D Wexner

Julie Ann M Van Koughnett, Steven D Wexner, Department of Colorectal Surgery, Cleveland Clinic Florida, Weston, FL 33331, United States

Author contributions: Both authors contributed to research, writing and revisions of manuscript.

Supported by Dr. Wexner is a consultant and receives consulting fees in the field of fecal incontinence from: Incontinence Devices, Inc; Mediri Therapeutics, Inc.; Medtronic Inc.; Renew Medical; Salix Pharmaceuticals

Correspondence to: Steven D Wexner, MD, PhD(Hon), FACS, FRCS, FRCS(Ed), Department of Colorectal Surgery, Cleveland Clinic Florida, 2950 Cleveland Clinic Boulevard, Weston, FL 33331, United States. wexners@ccf.org

Telephone: +1-954-6596020 Fax: +1-954-6596021

Received: July 30, 2013 Revised: October 7, 2013

Accepted: November 2, 2013

Published online: December 28, 2013

Abstract

The severity of fecal incontinence widely varies and can have dramatic devastating impacts on a person's life. Fecal incontinence is common, though it is often under-reported by patients. In addition to standard treatment options, new treatments have been developed during the past decade to attempt to effectively treat fecal incontinence with minimal morbidity. Non-operative treatments include dietary modifications, medications, and biofeedback therapy. Currently used surgical treatments include repair (sphincteroplasty), stimulation (sacral nerve stimulation or posterior tibial nerve stimulation), replacement (artificial bowel sphincter or muscle transposition) and diversion (stoma formation). Newer augmentation treatments such as radiofrequency energy delivery and injectable materials, are minimally invasive tools that may be good options before proceeding to surgery in some patients with mild fecal incontinence. In general, more invasive surgical treatments are now reserved for moderate to severe fecal incontinence. Functional and quality of life related outcomes, as well

as potential complications of the treatment must be considered and the treatment of fecal incontinence must be individualized to the patient. General indications, techniques, and outcomes profiles for the various treatments of fecal incontinence are discussed in detail. Choosing the most effective treatment for the individual patient is essential to achieve optimal outcomes in the treatment of fecal incontinence.

© 2013 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Fecal incontinence; Treatment; Sacral nerve stimulation; Sphincteroplasty; Artificial bowel Sphincter; Biofeedback

Core tip: An increasing number of treatment options for the management of fecal incontinence have been developed. In addition to traditional options such as sphincteroplasty and colostomy, non-surgical options such as biofeedback and dietary modification may be considered for mild incontinence. Injectable materials and radiofrequency energy delivery are two newer treatments for mild incontinence. Surgical options for moderate to severe incontinence include sacral nerve stimulation, artificial bowel sphincter implantation, muscle transposition, antegrade continence enemas, sphincteroplasty, and colostomy formation. Treatment for fecal incontinence (repair, stimulation, replacement, augmentation, or diversion) must be individualized to the patient, considering the underlying cause and impact on quality of life of the fecal incontinence.

Van Koughnett JAM, Wexner SD. Current management of fecal incontinence: Choosing amongst treatment options to optimize outcomes. *World J Gastroenterol* 2013; 19(48): 9216-9230 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v19/i48/9216.htm> DOI: <http://dx.doi.org/10.3748/wjg.v19.i48.9216>

INTRODUCTION

Fecal incontinence is a common problem; one that is likely underreported in the general population. The prevalence of fecal incontinence varies in the literature, with one study of over 4000 surveyed American adults finding a prevalence of 8.3%^[1]. The much larger and more recent Mature Women's Health Study of over 5800 American women found an even higher incidence of accidental bowel leakage of almost 20%^[2]. Incontinence to liquid or solid stool, mucous, or flatus occurs with varying frequency and can have a range of impact on daily function^[1]. The Mature Women's Health Study found that nearly 40% of women with accidental bowel leakage have severe symptoms impacting their quality of life, even though less than one third of women sought medical care for their bowel leakage^[3,4]. While there can be many etiologic factors contributing to its development, there are some common risk factors. Age, diarrhea or frequent bowel movements, nocturnal bowel movements, other bowel disorders, and the presence of urinary incontinence are commonly associated with fecal incontinence^[1,4,5]. In women, internal sphincter injury and reduced perineal descent related to obstetrical trauma independently predict the development of fecal incontinence^[6]. Other risk factors include neurological disorders, congenital anorectal malformations, trauma, iatrogenic injury during anorectal procedures, and chronic diseases such as diabetes^[6-9].

It is necessary to complete a physiological and anatomical assessment of the pelvis and colon in order to choose the most appropriate treatment option for a patient's fecal incontinence. This caveat is especially important since many women with fecal incontinence have associated genital and urinary anatomical or functional problems^[10]. A rectal examination may identify a sphincter defect or decreased rectal tone. This finding may be helpful to identify potential etiologies and treatments for a patient's fecal incontinence. Though not all investigations are required for every patient, options include anal or pelvic ultrasound, anal manometry, defecography, magnetic resonance imaging, and electromyography with pudendal nerve terminal motor latency testing. Anatomical imaging can help identify sphincter defects and associated pelvic floor disorders such as rectocele or prolapse, which may be contributing to the severity of incontinence^[11,12]. A physiology lab is helpful for the assessment of incontinence and other pelvic floor disorders.

The impact of fecal incontinence varies and can greatly alter a person's ability to perform daily activities. One may alter timing of meals or eating habits, and possibly avoid all social occasions for fear of embarrassment^[8]. While fecal incontinence is not a normal part of aging it may be perceived as such, and older people may not seek treatment until symptoms are severe. Treatment options for fecal incontinence range from dietary modification and physical therapy to major surgery, such as colostomy formation. In recent decades, many new treatments for fecal incontinence have been developed with

good success, adding to traditional options of sphincteroplasty and ostomy formation. These alternatives include biofeedback, radiofrequency, injectable materials, and surgical approaches such as sacral nerve stimulation, the artificial bowel sphincter, and muscle transposition. A recent Cochrane review concluded that there is insufficient evidence to allow for quality comparisons to be made among the various surgical approaches to fecal incontinence^[13]. The decision among these options is multifactorial and the severity of the incontinence, patient anatomy, and patient wishes must all be carefully considered. The aim of this article is to review current options for the management of fecal incontinence, their indications, and reported outcomes. The treatments most commonly offered by the authors, from the five available categories of repair, stimulation, replacement, augmentation, and diversion, are discussed.

DIETARY MODIFICATION AND MEDICATION

Modifiable diet and lifestyle factors may be identified which can provide simple interventions to try to improve symptoms. Smoking and sedentary lifestyle are associated with fecal incontinence^[14]. Weight loss has been shown to improve fecal incontinence in obese women^[15]. Medications should be reviewed with the help of a pharmacist to identify potentially incriminating medications. Low fiber and high fat diets may be contributory to loose stools. Loose stools and diarrhea often precipitate symptoms of fecal incontinence and may be improved with dietary and medication alterations. Other factors may be identified that may suggest the need for further testing or anatomical causes of fecal incontinence. For example, cholecystectomy may lead to persistent diarrhea and flatulence which may amplify symptoms of fecal incontinence; cholestyramine may help relieve these symptoms^[16,17].

The addition of a daily fiber supplement should be advocated in fecal incontinence. It acts as a bulking agent to allow for more solid stool and adds little to no morbidity to the patient. A randomized, blinded, placebo controlled study found that fiber improved fecal incontinence and stool consistency within 1 mo in the community living population^[18]. In addition to fiber, medications with a constipating effect may be useful for patients with fecal incontinence with loose stools. These pharmacologic agents include loperamide, diphenoxylate and atropine, and codeine. Loperamide is most commonly used and may also have beneficial effects on anal sphincter resting tone^[19]. Unfortunately, studies comparing various medications are lacking and trials of medications for the treatment of fecal incontinence include very heterogeneous populations and treatments^[20]. A Cochrane review conducted in 2013 concluded that there is insufficient evidence to guide the decision between medications for the treatment of incontinence in various clinical situations^[20]. Clearly, no medication will cure moderate to severe fecal incontinence, but it should certainly be utilized in mildly

Table 1 Success of biofeedback for fecal incontinence

Ref.	Year	Patients (n)	Significant reduction in incontinence (percentage of patients)	Improvement in quality of life (percentage of patients)	Adjuncts to traditional biofeedback
Keck <i>et al</i> ^[33]	1994	15	73%	NR	None
Solomon <i>et al</i> ^[28]	2003	102	70%	69%	Anal manometry, transanal ultrasound
Terra <i>et al</i> ^[34]	2006	239	60%	NR	EMG, electrostimulation
Naimy <i>et al</i> ^[30]	2007	49	None	None	Electrostimulation
Byrne <i>et al</i> ^[32]	2007	385	70%	87%	None
Heymen <i>et al</i> ^[27]	2009	45	76%	NR	None
Schwandner <i>et al</i> ^[29]	2010	158	50%	NR	EMG, electrostimulation
Bartlett <i>et al</i> ^[26]	2011	72	86%	100%	None
Jodorkovsky <i>et al</i> ^[31]	2013	12	80%	NR	None

NR: Not reported; EMG: Electromyography.

symptomatic patients where indicated.

BIOFEEDBACK

Biofeedback is a form of physical therapy and muscle re-training offered to patients refractory to medical treatment of fecal incontinence. There are numerous regimens, most of which involve many weeks of treatment lead by a physical therapist. Numerous studies have attempted to define the most effective regimen and most responsive patient population, but overall there are few high quality studies showing a definitive impact of biofeedback on fecal incontinence^[21]. It has been suggested by some authors that biofeedback should be offered to all patients who have not responded to medical interventions of fecal incontinence because it is safe, inexpensive, and effective long term^[22]. Older patients with normal defecation physiology appear to respond well^[23]. Advanced anorectal physiology tests such as manometry, defecography, pelvic magnetic resonance imaging, and pudendal nerve terminal motor latency testing do not seem to predict who will respond best to biofeedback^[24]. Patients with mild or moderate fecal incontinence who have not responded well to medical treatments are likely the best candidates for biofeedback^[25].

The technique of biofeedback may include monitored or home sessions, pelvic floor exercises, digital feedback, electrical stimulation, balloons, and manometric or ultrasound monitoring of response. Pelvic floor exercises alone have been shown to improve fecal incontinence scores and quality of life^[26]. In one study of pelvic floor exercises, no differences in treatment effect were found between the different regimens, but symptoms improved in both groups^[26]. The addition of biofeedback using manometry is more effective than pelvic floor exercises alone to improve fecal incontinence scores and achieve more physiologically normal defecation^[27]. Biofeedback with digital feedback alone may be just as effective as manometry and ultrasound guided treatment, providing enough feedback to guide re-training, as found in a randomized controlled trial of different methods of biofeedback^[28]. Some literature suggests that electrical stimulation leads to more effective results over biofeedback alone, while others have found that biofeedback alone is

adequate to improve patient symptoms^[29,30]. A multicenter randomized and blinded trial found that the combination of electrical stimulation with extended treatment duration (longer than 3 mo) achieved the best results^[29]. Such treatment regimens may not be available in many centers, but access to a trained biofeedback therapist who is aware of the various treatment modalities may be invaluable to the population with fecal incontinence.

Biofeedback requires the patient and therapist to commit to treatment for a number of weeks to months. One study found that only 44% of patients with fecal incontinence who were recommended to undergo biofeedback therapy completed the treatment^[31]. This finding was largely due to lack of insurance coverage and distance to treatment centers^[31]. It is important to note that in this study those patients who did undergo biofeedback reported an 80% positive response to the treatment^[31]. Other studies have confirmed improvement in over 70% of patients when fecal incontinence scores and quality of life scores were assessed^[32,33]. Table 1 summarizes the success of biofeedback. Physiologic parameters such as squeeze pressure and maximum tolerated volume have also been reported to improve with biofeedback^[34]. Improvements in fecal incontinence scores are durable over at least 1 year, but some patients may require additional sessions to boost the effect^[35]. Pelvic floor training with biofeedback is likely beneficial to many patients with fecal incontinence long term, but patients and therapist must be willing to devote the time to a complete set of sessions to see maximum benefit. In those able to do so, biofeedback may achieve improvement in symptoms without invasive procedures.

REPAIR

Sphincteroplasty

Sphincteroplasty has long been the standard of care of the management of fecal incontinence related to anal sphincter injury^[36]. The vast majority of patients who undergo sphincteroplasty have a history of vaginal delivery^[37]. However, only about one third of women who have had a known sphincter injury related to vaginal delivery develop fecal incontinence over time^[36]. Pudendal nerve injury, failed prior sphincteroplasty, multiple

Table 2 Success of overlapping sphincteroplasty

Ref.	Year	No. of patients with follow-up	Mean follow-up (mo)	Success ¹ (percentage of patients)
Karoui <i>et al</i> ^[52]	2000	74	40	28%
Halverson <i>et al</i> ^[40]	2002	49	69	46%
Bravo Gutierrez <i>et al</i> ^[39]	2004	130	120	41%
Barisic <i>et al</i> ^[49]	2006	65	80	48%
Maslekar <i>et al</i> ^[55]	2007	64	84	80%
Oom <i>et al</i> ^[50]	2009	120	111	60%
Mevik <i>et al</i> ^[51]	2009	25	84	53%
Zutshi <i>et al</i> ^[53]	2009	31	129	0%

¹Success variably defined in studies. Good, excellent or complete continence included as success.

vaginal deliveries, history of third or fourth degree tear, and instrument-assisted vaginal deliveries are all factors which may predispose to fecal incontinence associated with sphincter defect and impact the success of sphincteroplasty^[38]. It is important to note that the majority of recent studies indicate that pudendal nerve injury as demonstrated by prolonged pudendal nerve terminal motor latency does not independently predict the success of sphincteroplasty^[39-41]. Many women who undergo sphincteroplasty have associated pelvic floor injuries, which do not seem to impact the success of sphincteroplasty^[42]. In addition, the combination of internal and external anal sphincter defect repair can lead to successful and equivalent outcomes when compared to external anal sphincter defect repair, alone^[43].

While various techniques for sphincteroplasty have been described, the most commonly performed procedure is the anterior overlapping sphincteroplasty. A curvilinear incision is made on the perineum and dissection proceeds until the edges of the external anal sphincter are identified and isolated. Care is taken to not dissect too far laterally to avoid nerve injury. The ends are overlapped and sutured together, providing new bulk to the sphincter complex and an intact circumferential ring of sphincter. Separate attention to the imbrication of the internal anal sphincter does not seem to add to the overall durability of the sphincteroplasty if the internal sphincter is not injured^[44]. Post-operative manometry shows significant increases in the length of the high pressure zone and resting and squeeze pressures^[37]. A diverting stoma is not required to achieve optimal outcomes in early repair of third and fourth degree tears during vaginal delivery^[45]. Delayed repair is associated with higher overall cost in this situation, but may still achieve good long term outcomes and may be the safer option depending on the clinical scenario^[45,46].

Posterior sphincter repair is rarely needed, given that most sphincter injuries are associated with traumatic vaginal delivery. However, posterior repair may be occasionally utilized for neurogenic fecal incontinence, multifocal sphincter defects, or after failed anterior sphincteroplasty in order to avoid any significant scar tissue in the area.

A similar technique is used as in the anterior technique, with a curvilinear posterior incision being used for access to the external anal sphincter. Some surgeons may proceed with a combined anterior and postanal approach, though this combination is not common. The success rate of the postanal approach is likely equivalent or less durable compared to anterior sphincteroplasty^[47,48]. In the absence of a specific iatrogenic posterior sphincter injury or excessive anterior scar tissue, the anterior sphincteroplasty should be considered the preferred approach.

The long term functional outcomes following anal sphincteroplasty are not ideal. The Wexner fecal incontinence score is commonly used to assess for incontinence following sphincteroplasty. In the short term, good results are achieved in over 70% of patients and excellent results in over half of patients^[49]. However, the long term outcomes which have been reported in numerous retrospective studies reveal a consistent decrease to 15% to 60% good long term continence^[39,40,50-56]. Interestingly, there is poor correlation between long term quality of life scores and fecal incontinence scores, with one study reporting that 95% of patients were satisfied with their operation a mean of 7 years following sphincteroplasty^[39,53]. A summary of long term outcomes is found in Table 2. Age has long been felt to be a predictor of success of sphincteroplasty, with many studies reporting that older patients do not have as durable long term outcomes compared to younger patients^[39,53,56]. However, a recent large review of 321 women who underwent sphincteroplasty showed that age is not a predictor of long term incontinence scores^[57]. A review of both sphincteroplasty and sacral nerve stimulation concluded that sphincteroplasty remains a good option for the management of incontinence due to sphincter defect, despite new technologies^[58]. Patients must be chosen after appropriate pre-operative evaluation to achieve optimal outcomes.

STIMULATION

Sacral nerve stimulation

For many patients and practitioners, sacral nerve stimulation has revolutionized the treatment of moderate to severe fecal incontinence. Adapted from its use in urinary incontinence, it may provide effective relief from fecal incontinence without any direct intervention on the anal sphincter complex. Interestingly, one study found that the only positive predictors of successful treatment with sacral nerve stimulation were loose stools and low stimulation intensity during the test phase of the procedure^[59]. Conversely, age, gender, etiology of fecal incontinence, and physiology study results did not impact the efficacy of sacral nerve stimulation^[59]. Though sacral nerve stimulation and sphincteroplasty have not been directly compared in the literature, numerous studies have shown that patients with sphincter defects can have excellent results with sacral nerve stimulation^[60-64]. The success of sacral nerve stimulation in these patients also does not appear to be correlated to the degree of sphincter defect^[63]. Pa-

Table 3 Studies of outcomes of sacral nerve stimulation

Ref.	Year	Patients (n)	Significant reduction in incontinence scores and incontinent episodes	Significant increase in quality of life
Leroi <i>et al</i> ^[73]	2005	27	Y	Y
Boyle <i>et al</i> ^[63]	2009	15	Y	NR
Brouwer <i>et al</i> ^[64]	2010	55	Y	Y
Wexner <i>et al</i> ^[79]	2010	120	Y	Y
Hollingshead <i>et al</i> ^[76]	2011	18	Y	NR
Lim <i>et al</i> ^[78]	2011	41	Y	Y
Mellgren <i>et al</i> ^[81]	2011	83	Y	Y
George <i>et al</i> ^[77]	2012	23	Y	Y
Devroede <i>et al</i> ^[83]	2012	78	Y	Y
Hull <i>et al</i> ^[75]	2013	76	Y	Y
Damon <i>et al</i> ^[82]	2013	92	Y	Y

Y: Yes; NR: Not reported.

tients known to have pudendal nerve injuries or previous sphincteroplasty can have good responses to sacral nerve stimulation^[64].

The mechanism by which sacral nerve stimulation improves fecal incontinence is not well defined, as it is multifactorial. A systematic review found that sacral nerve stimulation likely works in 3 ways: stimulation of a somato-visceral reflex, direct effect on the anal sphincter complex, and afferent nerve modulation^[65]. It is postulated that sacral nerve stimulation may induce a change in anal sphincter muscle type from fast to slow twitch, thus reducing muscle fatigue, though this has not been definitively demonstrated in the sacral nerve stimulation population^[66]. Sensory changes include the sensation of rectal filling and urge to defecate at higher rectal volume^[67]. Sacral nerve stimulation alters colonic transit by inducing retrograde colonic propagating sequences, activity which may slow transit in the setting of fecal incontinence^[68]. In an animal model, sacral nerve stimulation was found to increase activity in the central cerebral cortex^[69]. The effects of sacral nerve stimulation are well beyond local effect on the anal sphincter complex.

There are two approaches to the implantation of the sacral nerve stimulator. Some surgeons introduce a peripheral nerve stimulator wire in the office, guided by anatomical landmarks. The patient is tested for response for a period of 1-2 wk and if good response is achieved, the permanent tined lead and stimulator device are implanted in the same setting in the operating room. The authors' preferred approach is a two-stage operative technique. The first stage is the insertion of the tined lead into the S3 foramen in the operating room with careful fluoroscopic and patient-directed guidance. Local anesthetic injections and light sedation allow the patient to signal when stimulation is felt in the perianal, perineal, or saddle regions during lead electrostimulation. In addition, sphincter bellows and plantar flexion of the great toe on the side of lead placement are used to further indication stimulation of the sacral nerve. Once a good response

is achieved the lead is tunneled into position. A temporary device is used during a 2 wk test phase. If a good response is achieved during the test phase, the patient undergoes a second procedure to implant the permanent device which is attached to the tined lead. This approach is associated with very little lead migration during the test phase but does require two operations. A test phase is important in both approaches, as not all patients will have a good response to lead placement^[70]. Each permanent device is programmed to the individual's response pattern. Successful strategies to prolong the durability of the device battery beyond the average of six years include cyclical stimulation and subsensory stimulation^[71,72].

Results of the first randomized multi-center study of sacral nerve stimulation were reported in 2005, showing that fecal incontinence was improved when the sacral nerve stimulator was activated^[73]. Longer term results are now available. Compared to medical treatment of fecal incontinence, sacral nerve stimulation is significantly more effective^[74]. A recent report from the SNS Study Group showed that in patients followed for at least 5 years, 89% have significant continued reduction in fecal incontinence and 36% had a complete response to sacral nerve stimulation^[75]. Numerous other studies from around the world have demonstrated significant long term reduction in fecal incontinence scores^[75-79]. Table 3 summarizes the results of studies of outcomes of sacral nerve stimulation. Furthermore, in women who have undergone sacral nerve stimulation for fecal incontinence; urinary, sexual, and vaginal symptoms also improve with a global benefit on pelvic floor health^[80]. Quality of life scores are also improved in the short and long term after sacral nerve stimulation^[79,81-84].

There are potential morbidities with sacral nerve stimulation including a 5% risk of lead displacement associated with the percutaneous lead testing technique^[85]. Pain at the surgical site and paresthesias are the most commonly reported complaints^[81]. Infection of the permanent device or surgical site occurs in 10%, with about half of those infections requiring surgical management^[81,85]. Overall, about one third of patients required surgical manipulation of the device in a study of long term outcomes^[75]. Despite potential morbidity associated with the device, sacral nerve stimulation has been shown to be cost-effective in the treatment of fecal incontinence^[86,87]. When balancing the effectiveness, morbidity profile, and cost-effectiveness of the technique, sacral nerve stimulation is a very valuable tool for the treatment of fecal incontinence, especially in its more severe forms.

REPLACEMENT

Artificial bowel sphincter

The artificial bowel sphincter is considered only for patients with severe fecal incontinence. It is an effective device, but requires long term follow up and a motivated patient. The use of an artificial bowel sphincter requires both manual dexterity and mental capacity to operate the device^[88]. Due to the high incidence of adverse events,

Table 4 Outcomes of artificial bowel sphincter

Ref.	Year	Patients (n)	Explanted devices (n)	Success (percentage of patients), intention to treat	Complications
Lehur <i>et al</i> ^[93]	2000	24	7	83%	Obstructed defecation
Altomare <i>et al</i> ^[94]	2001	28	3	75%	Obstructed defecation, infection, device erosion
Devesa <i>et al</i> ^[95]	2002	53	10	65%	Perforation, infection, sepsis, device erosion, pain, impaction
Wong <i>et al</i> ^[97]	2002	112	41	53%	Infection, pain
Lehur <i>et al</i> ^[101]	2002	16	4	69%	Erosion
Parker <i>et al</i> ^[96]	2003	45	18	49%	Infection, pain
O'Brien <i>et al</i> ^[98]	2004	14	1	NR as percentage	Obstructed defecation, non-healing of wound
Melenhorst <i>et al</i> ^[103]	2008	33	7	NR as percentage	Pain, perforation, infection, obstructed defecation
Ruiz Carmona <i>et al</i> ^[99]	2009	17	11	53%	Infection, erosion
Wexner <i>et al</i> ^[90]	2009	51	31	NR as percentage	Infection, malfunction, erosion, pain
Wong <i>et al</i> ^[100]	2011	52	14	67%	Perforation, cuff leak

NR: Not reported.

other treatment options should be considered and attempted before proceeding to artificial bowel sphincter^[89]. Contraindications include Crohn's disease, local sepsis, prior radiation, poor quality of the perineal tissues, severe constipation, and incontinence associated irritable bowel syndrome^[89]. Disruption of the anal sphincter complex due to trauma, severe obstetrical injury, and imperforate anus are common indications^[89,90]. Sacral nerve stimulation and the artificial bowel sphincter have largely replaced muscle transposition and dynamic graciloplasty for the treatment of severe fecal incontinence, with better functional outcomes and quality of life parameters^[91,92]. Patients must be carefully selected and extensively counselled on the risks and benefits of the artificial bowel sphincter, as discussed below.

Meticulous sterile technique and thorough bowel preparation are essential to reduce the risk of infection associated with the artificial bowel sphincter. The 3 components of the artificial bowel sphincter are connected *via* tubing and compose the sphincter cuff, the reservoir balloon, and control pump. These components are inserted *via* perineal, Pfannenstiel, and labial or scrotal incisions, respectively. The cuff itself is chosen for size based on circumferential length around the rectum and width. It is inserted first and great care is taken to ensure there is adequate tissue bulk distal to the cuff, in an attempt to avoid device erosion and infection. The balloon holds approximately 40 mL of liquid and is left filled with the device deflated at the end of the procedure after testing the control pump. The device is not activated for four to six weeks to allow for complete healing. The patient is taught how to fill and empty the cuff by using the implanted control pump.

Patients who retain the artificial bowel sphincter long term have reported very good functional and qualitative results. Manometry results show that the artificial bowel sphincter achieves normal resting tone when the cuff is filled^[93]. Improved continence is achieved in over 75% of patients, with one series reporting normal continence in two-thirds of patients^[94,95]. Though adverse events are

significant, patients who retain the device have excellent responses to artificial bowel sphincter implantation based on incontinence scores^[93-100]. Quality of life scores are also markedly improved after successful treatment of fecal incontinence with the artificial bowel sphincter^[96,98,99,101]. A systematic review of the safety of the artificial bowel sphincter noted that functional outcomes and quality of life scores for those patients who do not retain a functioning device are not reported in the literature^[102].

Complications following artificial bowel sphincter implantation unfortunately remain high and often lead to device explantation, mitigating the overall population benefit in fecal incontinence. Unfortunately, these complications continue to accrue long term^[90]. The rate of revision of the device has been reported to be up to 50%, with infection and device failure the most common reasons^[100]. About 25%-40% of artificial bowel sphincters become infected over time^[90,100,103]. Erosion of the cuff or control pump and post-operative constipation may also occur^[92,104,105]. The outcomes and complications associated with the artificial bowel sphincter are included in Table 4. In summary, a balanced consideration of potential benefits and adverse events is important and artificial bowel sphincter may still be the optimal treatment consideration for select patients with severe fecal incontinence.

Muscle transposition

Muscle transposition is a technique used to physically replace the sphincter with *in vivo* muscle bulk. It is most often used in the setting of a traumatic or iatrogenic disruption of the anal sphincters to recreate a wrap of muscle around the anus. A substantial congenital or post-traumatic defect is indicated to consider muscle transposition. The two muscles widely described in the literature for transposition are the gluteus maximus and gracilis muscles. These are useful because of their proximity to the anus, sizeable muscle bulk, and nerve locations which are amenable to preservation upon transposition. In addition, the gluteus maximus was thought to be a good

Table 5 Outcomes of graciloplasty

Ref.	Year	Type of graciloplasty	Patients (n)	Success (percentage of patients)
Kumar <i>et al</i> ^[114]	1995	Unstimulated	9	100%
Eccersley <i>et al</i> ^[113]	1999	Unstimulated	8	100%
Madoff <i>et al</i> ^[109]	1999	Stimulated	128	66%
Wexner <i>et al</i> ^[110]	2002	Stimulated	115	62%
Bresler <i>et al</i> ^[112]	2002	Stimulated	24	79%
Rongen <i>et al</i> ^[111]	2003	Stimulated	200	72%
Thornton <i>et al</i> ^[117]	2004	Stimulated	38	73%
Hassan <i>et al</i> ^[107]	2010	Stimulated	31	71%

choice for transposition given that involuntary gluteal contraction occurs with the strong urge to avoid involuntary defecation^[106].

The surgical technique of muscle transposition is complex and requires significant experience to gain expertise. Three main options exist: gluteoplasty, graciloplasty, and dynamic (or stimulated) graciloplasty. Gluteoplasty is performed with the patient in the prone position with the table flexed at the hips. Bilateral incisions over the gluteus are made and two tongues (one from each side) of the lower 10% of the muscle are raised with care taken to preserve the neurovascular bundles^[106]. The mobilized muscle is then tunnelled and delivered through separate bilateral curvilinear incisions around the anus. The contralateral mobilized segments are sutured together to create a ring of muscle.

In a graciloplasty procedure, the patient is placed in the modified lithotomy position. Two or three incisions are made along the longitudinal access of the gracilis muscle on the chosen side to harvest the entire length of the gracilis. The neurovascular bundle is preserved through its identification during medial dissection. The muscle is released distally and tunneled medially. A perineal incision is made and the gracilis is wrapped circumferentially around the anus. In the dynamic graciloplasty technique, an electrode is placed in the gracilis muscle and an implantable device similar to that used for sacral nerve stimulation is implanted in the abdominal wall. Modified approaches to dynamic graciloplasty include temporary stimulation with an external stimulator for muscle retraining, similar to biofeedback^[107]. It must be noted that the stimulator and leads for dynamic graciloplasty are not currently approved for use in North America.

Much like the artificial bowel sphincter, muscle transposition has fairly good functional outcomes but high rates of complications and re-operation; graciloplasty has largely replaced gluteoplasty. The largest and most recent study of gluteoplasty reported a good functional outcome in 59% of patients^[108]. Successful functional outcomes for graciloplasty, dynamic and unstimulated, is consistently reported to be about 60%-75%, with earlier success of unstimulated graciloplasty being even higher^[107-114]. Table 5 lists the published success rates of graciloplasty. If a patient has a stoma at the time of the graciloplasty, eventual outcomes are equivalent to those

who do not have a stoma, but are delayed in achieving them^[110]. Complications of the procedure are common, and include surgical site infections, pain, rectal injury, and erosion of the device in the case of dynamic graciloplasty^[112,115,116]. In addition, constipation due to obstructed defecation is commonly reported in as many as 50% of patients^[115-117]. There are no studies directly comparing muscle transfer to other surgical treatments of fecal incontinence. Graciloplasty followed by artificial bowel sphincter implantation may be the best combination option for adult patients with fecal incontinence attributable to congenital imperforate anus^[118].

DIVERSION

Antegrade continence enema

The antegrade continence enema was first described by Malone *et al*^[119] in 1990. It is used to control fecal soiling in both adults and children, but is most commonly used and reported in the pediatric population. Neurogenic conditions, such as spina bifida, resulting in neurogenic bowel and urinary symptoms are the most common indications in children. While the antegrade continence enema may be helpful in pure fecal incontinence, most often patients who undergo this procedure have the combination of constipation or colonic dysmotility with associated overflow fecal incontinence. Patients also commonly undergo urological procedures at the same time to control neurogenic bladder symptoms, with good results for these combined indications^[120]. In adults, good functional outcomes are better in this setting, when compared to those patients who undergo the procedure for constipation alone^[121]. While an antegrade continence enema does not alter anorectal physiology or anatomy, it provides a mechanism to empty the colon in a controlled fashion, allowing the patient to perform their daily activities with little worry of fecal soiling or incontinent episodes.

Since Malone's original description, various techniques have been described for the creation of an antegrade continence enema. The appendix, ileum, cecum, and left colon may be used successfully as the access point for irrigation^[122-124]. The appendix is most commonly used, where it is inverted and fixated to the skin at the umbilicus or right lower quadrant. This can be performed open or laparoscopically with good results^[124]. The access point is left intubated with a catheter for about 3 wk after the operation before intermittent intubations begin. Patients or their caregivers then intubate the bowel daily to every few days and perform colonic irrigation with tap water or an electrolyte or bowel cleansing solution. Both tap water and commercial products have good irrigation results, with solution irrigants achieving slightly better continence rates^[125]. The volume of irrigation is gradually increased over time after the procedure and the timing and frequency of irrigation through the site may be largely patient directed. In the pediatric patient population, the operation is performed around the age of 10 years.

Few studies report on outcomes of antegrade con-

Table 6 Outcomes of antegrade continence enema in incontinent adults

Ref.	Year	Patients using antegrade continence enema on follow-up	Percentage of patients achieving continence	Complication rate
Gerharz <i>et al</i> ^[121]	1997	8	100%	44%
Teichman <i>et al</i> ^[120]	1998	7	86%	71%
Teichman <i>et al</i> ^[128]	2003	4	75%	67%
Lefevre <i>et al</i> ^[127]	2006	18	94%	33%
Poirier <i>et al</i> ^[126]	2007	14	78%	67%

tinence enemas in adults. Overall, functional results are very good, with about 75% of adults achieving continence with the procedure^[126-128]. Quality of life improves in adult patients with antegrade continence enemas, although not all patients continue to use their antegrade continence enema in the long term^[127,128]. See Table 6 for a summary of antegrade continence enema study results. In children, full continence is achieved in 65%-100% of patients^[122,125,129-133]. Even though the amount of time devoted to bowel care may not significantly change, satisfaction and quality of life scores improve for most children and parents^[131,134-137]. Persistent leakage, stoma stenosis, and surgical site infections are common complications, with one study quoting a 13% chance of requiring stoma revision due to stoma complications^[130,131,138]. While the antegrade continence enema is not commonly performed in adults, the patients who have grown to adulthood require long term follow up and attention to these possible complications.

Fecal diversion

The creation of a colostomy or ileostomy provides definitive control of fecal incontinence. An ileostomy may be considered in patients with colonic transit abnormalities but the colostomy is the standard ostomy utilized in the treatment of fecal incontinence. In many patients the ostomy can be created using a laparoscopic approach to improve recovery time. While a colostomy is not without short and long term risks, such as bleeding, anesthesia related cardiac or respiratory morbidities, and parastomal hernia, it is a safe and effective treatment of severe fecal incontinence. It is generally only offered if other treatment modalities have failed. Patients are usually understandably very resistant to the idea of a permanent colostomy, fearing it will be difficult to manage and have great impact on self-image and social interactions.

When patients who had undergone colostomy creation for fecal incontinence were surveyed, general quality of life and fecal incontinence quality of life scores were actually higher in the colostomy group when compared to other patients with fecal incontinence^[139]. Another study found that patients generally reported high satisfaction levels with their stomas for fecal incontinence, with over 80% of patients stating that they would likely or definitely choose to undergo the procedure again^[140]. Compared to other surgical treatments of sever incontinence (dynamic

Table 7 Outcomes of radiofrequency energy treatments

Ref.	Year	Patients (n)	Significant improvement in incontinence scores after treatment	Significant improvement in quality of life
Efron <i>et al</i> ^[143]	2003	50	Y	Y
Felt-Bersma <i>et al</i> ^[147]	2007	11	Y	NR
Takahashi-Monroy <i>et al</i> ^[142]	2008	19	Y	Y
Lefebure <i>et al</i> ^[144]	2008	15	Y	N
Kim <i>et al</i> ^[148]	2009	8	N	N
Ruiz <i>et al</i> ^[145]	2010	24	Y	Y
Abbas <i>et al</i> ^[146]	2012	27	Y	NR

Y: Yes; N: No; NR: Not reported.

graciloplasty and artificial bowel sphincters), a British study found colostomy to be most cost effective in terms of quality adjusted life years^[92]. While fecal diversion is not required in the majority of patients presenting for treatment of fecal incontinence, it is a viable, definitive, and well-tolerated treatment which offers good quality of life.

AUGMENTATION

Radiofrequency energy

There is a gap between medical and surgical treatment options in fecal incontinence^[141]. Radiofrequency energy delivery and injectable materials are becoming increasingly popular as minimally invasive procedural treatments that may bridge this gap. The delivery of radiofrequency energy to the internal anal sphincter, known as the SECCA[®] procedure, is proposed to induce local restructuring of collagen, leading to a more robust internal anal sphincter and better continence. It can be used for patients with mild or moderate fecal incontinence who are unwilling or not candidates to undergo surgical treatment after failing medical management. It may also be applied to patients with idiopathic or sphincter defect-associated fecal incontinence.

The technique of radiofrequency energy delivery is simple. It is done with conscious sedation and local anesthesia on an outpatient basis in endoscopy or the operating room. A commercial device is utilized and the procedure takes about 30 min. The device resembles a clear plastic anoscope with four retractable needles. The needles are electrodes which are deployed into the anorectal mucosa to deliver radiofrequency energy to the internal anal sphincter, starting just distal to the dentate line and moving proximally. The device delivers radiofrequency while simultaneously monitoring the temperature and impedance of the tissues to avoid burning. The device is activated four or five times per quadrant of the anorectum, moving 5 mm more proximal before each activation in a quadrant. The machine provides constant feedback on the contact with the tissues, temperature and impedance during the device activation, and the timing of each activation, giving visual and sound cues to the surgeon

Table 8 Outcomes of dextranomer in hyaluronic acid gel for fecal incontinence

Ref.	Year	Patients (n)	> 50% reduction in fecal incontinence episodes (percentage of patients)	Significant quality of life improvement
Dodi <i>et al</i> ^[150]	2010	115	64%	Yes
Graf <i>et al</i> ^[161]	2011	136	52%	Yes
Schwandner <i>et al</i> ^[159]	2011	21	56%	Yes
Danielson <i>et al</i> ^[160]	2012	34	76%	Yes
La Torre <i>et al</i> ^[156]	2013	83	63%	Yes

throughout the procedure.

Reports of the success of radiofrequency energy treatment are generally, though not universally, positive and are summarized in Table 7. Numerous studies have reported long term improvement in fecal incontinence scores^[142-145]. The cohort with the longest reported follow-up showed a durable reduction in mean Wexner fecal incontinence scores from 14 to 8 and found that most participants had a greater than 50% improvement in symptoms after 5 years^[142]. Similarly, patient satisfaction and quality of life scores show improvement after radiofrequency energy treatment^[142-145]. Another study with a higher average baseline fecal incontinence score compared to other trials found that only 22% of patients had sustained treatment benefits at an average follow up of 40 mo^[146]. Despite the overall favorable outcomes of radiofrequency energy delivery, anal manometry testing does not show any significant change in physiologic parameters^[143,147,148]. No major adverse events have been reported following radiofrequency energy delivery, though there have been reports of infection, hematoma, minor bleeding, and anal pain^[145,147,148].

Injectable materials

Various injectable materials have included trialed for local injection of the sphincter complex to treat fecal incontinence. Benefits of this approach are that it is an outpatient procedure with little discomfort that has low morbidity. The materials used have included collagen, silicone, autologous fat, glutaraldehyde, carbon-coated beads, dextranomer in hyaluronic acid gel, and others^[149]. Dextranomer in hyaluronic acid gel (NASA/Dx) has received the most extensive recent investigation and attention in the literature. Injectables may be used in patients who have failed medical treatment and have fecal leakage or mild to moderate fecal incontinence^[149]. The bulking effect may not be permanent and may require repeat injections at subsequent office visits.

The technique of injection is relatively simple. The open-label multicenter trial of NASA/Dx involved four quadrant injections of 1 mL of NASA/Dx into the deep submucosa of the anal canal^[148]. This was performed through an anoscope and done with the patient in the prone jack-knife or lithotomy positions. The injections were placed at a 30 degree angle 5-10 mm proximal to the dentate line^[150]. The needle was kept in place for

up to 30 s so that the gel would not leak from the site^[150]. There are very few comparative trials amongst injectable materials. A small study of 40 patients found that silicone was more effective than carbon-coated beads to reduce incontinence^[151]. No published studies have compared NASA/Dx with other injectables. One randomized controlled trials comparing NASA/Dx to biofeedback and found no significant difference in functional outcomes^[152]. Biofeedback, however, certainly requires more dedication and long term commitment from the patient. The effect of injectables on manometry parameters are an increase in the length of the high pressure zone and asymmetry index^[153]. The impact on resting pressure is variable in the literature, ranging from improvements in resting pressure to no effect^[153,154].

There are no long term outcomes reported yet for NASA/Dx, the most popular injectable. The longest reported outcomes are at 2 years^[155,156]. A Cochrane review published in 2013 noted the absence of long term studies, making definitive conclusions about the utility of injectables difficult^[157]. See Table 8 for a summary of cohort studies investigating the utility of NASA/Dx gel. A good response is considered a 50% reduction in the number of reported incontinence episodes, which is reported to occur in over 50% of patients who have been treated with injectables^[150,154,156,158-161]. In addition, the majority of patients have good quality of life improvement, as reported on both global quality of life and fecal incontinence quality of life scores^[150,155,156,158]. Morbidity from the use of injectables is low, with fever and proctalgia being the two most common adverse events and bleeding, abscess, and pain being other rare reported events^[150,156,160,161]. Though many patients with fecal incontinence may be candidates for the use of injectables, the ideal candidate is one who has seepage or mild to moderate incontinence who has failed medical management but is not yet ready to pursue surgical treatment. Prior use of an injectable such as NASA/Dx does not preclude future surgical treatments such as sacral nerve stimulation, sphincteroplasty or artificial bowel sphincter.

CONCLUSION

Successful treatment of fecal incontinence requires careful consideration of the individual patient's severity of incontinence. Treatments range from inexpensive medications and physical therapy to complex surgical procedures such as artificial bowel sphincter implantation and muscle transposition. In general, more invasive treatments are required for more severe incontinence or after less invasive treatments have failed. A careful history including obtaining an incontinence score, physical examination, bowel diary, and adjunctive anal physiology tests should be utilized to define the nature of the fecal incontinence. Minimally invasive approaches including biofeedback, radiofrequency energy, and injectables have moderate long term success. Sphincteroplasty remains an acceptable option for patients with documented sphincter defects. Be-

cause initially adequate functional outcomes decline over time, quality of life improvement after sphincteroplasty is not robust long-term. Sacral nerve stimulation is very effective in managing moderate to severe fecal incontinence and has had a great impact on the treatment of fecal incontinence. In the very long-term, patients will require additional procedures to change the battery of the sacral nerve stimulator but the procedure has excellent reproducible long term functional and quality of life outcomes. The artificial bowel sphincter has similar outcomes in those patients who retain the device, but further studies aimed at reducing infection, erosion, and device failure must be undertaken. Fecal diversion remains a good option for severe fecal incontinence and actually provides the patient with satisfying quality of life. Knowledge of these currently used treatments is essential to honest and thorough counseling of the patient with fecal incontinence to improve treatment success. Together with the patient, the surgeon can then best select treatment from the five available categories of repair, replacement, augmentation, stimulation, and diversion.

REFERENCES

- 1 **Whitehead WE**, Borrud L, Goode PS, Meikle S, Mueller ER, Tuteja A, Weidner A, Weinstein M, Ye W. Fecal incontinence in US adults: epidemiology and risk factors. *Gastroenterology* 2009; **137**: 512-517 [PMID: 19410574 DOI: 10.1053/j.gastro.2009.04.054]
- 2 **Brown HW**, Wexner SD, Segall MM, Brezoczky KL, Lukacz ES. Accidental bowel leakage in the mature women's health study: prevalence and predictors. *Int J Clin Pract* 2012; **66**: 1101-1108 [PMID: 23067034 DOI: 10.1111/ijcp.12018]
- 3 **Brown HW**, Wexner SD, Lukacz ES. Factors associated with care seeking among women with accidental bowel leakage. *Female Pelvic Med Reconstr Surg* 2013; **19**: 66-71 [PMID: 23442502 DOI: 10.1097/SPV.0b013e31828016d3]
- 4 **Brown HW**, Wexner SD, Segall MM, Brezoczky KL, Lukacz ES. Quality of life impact in women with accidental bowel leakage. *Int J Clin Pract* 2012; **66**: 1109-1116 [PMID: 23067035 DOI: 10.1111/ijcp.12017]
- 5 **Kang HW**, Jung HK, Kwon KJ, Song EM, Choi JY, Kim SE, Shim KN, Jung SA. Prevalence and predictive factors of fecal incontinence. *J Neurogastroenterol Motil* 2012; **18**: 86-93 [PMID: 22323992 DOI: 10.5056/jnm.2012.18.1.86]
- 6 **Bharucha AE**, Fletcher JG, Melton LJ, Zinsmeister AR. Obstetric trauma, pelvic floor injury and fecal incontinence: a population-based case-control study. *Am J Gastroenterol* 2012; **107**: 902-911 [PMID: 22415196 DOI: 10.1038/ajg.2012.45]
- 7 **Hayden DM**, Weiss EG. Fecal incontinence: etiology, evaluation, and treatment. *Clin Colon Rectal Surg* 2011; **24**: 64-70 [PMID: 22379407 DOI: 10.1055/s-0031-1272825]
- 8 **Hull T**. Fecal incontinence. *Clin Colon Rectal Surg* 2007; **20**: 118-124 [PMID: 20011386 DOI: 10.1055/s-2007-977490]
- 9 **Nelson RL**. Epidemiology of fecal incontinence. *Gastroenterology* 2004; **126**: S3-S7 [PMID: 14978632 DOI: 10.1053/j.gastro.2003.10.010]
- 10 **Maglinte DD**, Kelvin FM, Fitzgerald K, Hale DS, Benson JT. Association of compartment defects in pelvic floor dysfunction. *AJR Am J Roentgenol* 1999; **172**: 439-444 [PMID: 9930799 DOI: 10.2214/ajr.172.2.9930799]
- 11 **Kelvin FM**, Maglinte DD, Benson JT. Evacuation proctography (defecography): an aid to the investigation of pelvic floor disorders. *Obstet Gynecol* 1994; **83**: 307-314 [PMID: 8290201]
- 12 **Maglinte DD**, Hale DS, Sandrasegaran K. Comparison between dynamic cystocolpoproctography and dynamic pelvic floor MRI: pros and cons: which is the "functional" examination for anorectal and pelvic floor dysfunction? *Abdom Imaging* 2013; **38**: 952-973 [PMID: 22446896 DOI: 10.1007/s00261-012-9870-x]
- 13 **Brown SR**, Wadhawan H, Nelson RL. Surgery for faecal incontinence in adults. *Cochrane Database Syst Rev* 2010; (9): CD001757 [PMID: 20824829 DOI: 10.1002/14651858.CD001757.pub3]
- 14 **Townsend MK**, Matthews CA, Whitehead WE, Grodstein F. Risk factors for fecal incontinence in older women. *Am J Gastroenterol* 2013; **108**: 113-119 [PMID: 23090350 DOI: 10.1038/ajg.2012.364]
- 15 **Markland AD**, Richter HE, Burgio KL, Myers DL, Hernandez AL, Subak LL. Weight loss improves fecal incontinence severity in overweight and obese women with urinary incontinence. *Int Urogynecol J* 2011; **22**: 1151-1157 [PMID: 21567259 DOI: 10.1007/s00192-011-1444-x]
- 16 **Lamberts MP**, Lugtenberg M, Rovers MM, Roukema AJ, Drenth JP, Westert GP, van Laarhoven CJ. Persistent and de novo symptoms after cholecystectomy: a systematic review of cholecystectomy effectiveness. *Surg Endosc* 2013; **27**: 709-718 [PMID: 23052498 DOI: 10.1007/s00464-012-2516-9]
- 17 **Bharucha AE**, Zinsmeister AR, Schleck CD, Melton LJ. Bowel disturbances are the most important risk factors for late onset fecal incontinence: a population-based case-control study in women. *Gastroenterology* 2010; **139**: 1559-1566 [PMID: 20708007 DOI: 10.1053/j.gastro.2010.07.056]
- 18 **Bliss DZ**, Jung HJ, Savik K, Lowry A, LeMoine M, Jensen L, Werner C, Schaffer K. Supplementation with dietary fiber improves fecal incontinence. *Nurs Res* 2001; **50**: 203-213 [PMID: 11480529]
- 19 **Hallgren T**, Fasth S, Delbro DS, Nordgren S, Oresland T, Hultén L. Loperamide improves anal sphincter function and continence after restorative proctocolectomy. *Dig Dis Sci* 1994; **39**: 2612-2618 [PMID: 7995187 DOI: 10.1007/BF02087698]
- 20 **Omar MI**, Alexander CE. Drug treatment for faecal incontinence in adults. *Cochrane Database Syst Rev* 2013; **6**: CD002116 [PMID: 23757096 DOI: 10.1002/14651858.CD002116.pub2]
- 21 **Norton C**, Cody JD. Biofeedback and/or sphincter exercises for the treatment of faecal incontinence in adults. *Cochrane Database Syst Rev* 2012; **7**: CD002111 [PMID: 22786479 DOI: 10.1002/14651858.CD002111.pub3]
- 22 **Ozturk R**, Niazi S, Stessman M, Rao SS. Long-term outcome and objective changes of anorectal function after biofeedback therapy for faecal incontinence. *Aliment Pharmacol Ther* 2004; **20**: 667-674 [PMID: 15352915 DOI: 10.1111/j.1365-2036.2004.02125.x]
- 23 **Fernández-Fraga X**, Azpiroz F, Aparici A, Casaus M, Malagelada JR. Predictors of response to biofeedback treatment in anal incontinence. *Dis Colon Rectum* 2003; **46**: 1218-1225 [PMID: 12972966 DOI: 10.1007/s10350-004-6718-7]
- 24 **Terra MP**, Deutekom M, Dobben AC, Baeten CG, Janssen LW, Boeckxstaens GE, Engel AF, Felt-Bersma RJ, Slors JF, Gerhards MF, Bijnen AB, Everhardt E, Schouten WR, Berghmans B, Bossuyt PM, Stoker J. Can the outcome of pelvic-floor rehabilitation in patients with fecal incontinence be predicted? *Int J Colorectal Dis* 2008; **23**: 503-511 [PMID: 18228027 DOI: 10.1007/s00384-008-0438-8]
- 25 **Boselli AS**, Pinna F, Cecchini S, Costi R, Marchesi F, Violi V, Sarli L, Roncoroni L. Biofeedback therapy plus anal electrostimulation for fecal incontinence: prognostic factors and effects on anorectal physiology. *World J Surg* 2010; **34**: 815-821 [PMID: 20108095 DOI: 10.1007/s00268-010-0392-9]
- 26 **Bartlett L**, Sloots K, Nowak M, Ho YH. Biofeedback for fecal incontinence: a randomized study comparing exercise regimens. *Dis Colon Rectum* 2011; **54**: 846-856 [PMID: 21654252 DOI: 10.1007/DCR.0b013e3182148fe]
- 27 **Heymen S**, Scarlett Y, Jones K, Ringel Y, Drossman D, Whitehead WE. Randomized controlled trial shows biofeedback

- to be superior to pelvic floor exercises for fecal incontinence. *Dis Colon Rectum* 2009; **52**: 1730-1737 [PMID: 19966605 DOI: 10.1007/DCR.0b013e3181b55455]
- 28 **Solomon MJ**, Pager CK, Rex J, Roberts R, Manning J. Randomized, controlled trial of biofeedback with anal manometry, transanal ultrasound, or pelvic floor retraining with digital guidance alone in the treatment of mild to moderate fecal incontinence. *Dis Colon Rectum* 2003; **46**: 703-710 [PMID: 12794569 DOI: 10.1007/s10350-004-6643-9]
- 29 **Schwandner T**, König IR, Heimerl T, Kierer W, Roblick M, Bouchard R, Unglaube T, Holch P, Ziegler A, Kolbert G. Triple target treatment (3T) is more effective than biofeedback alone for anal incontinence: the 3T-AI study. *Dis Colon Rectum* 2010; **53**: 1007-1016 [PMID: 20551752 DOI: 10.1007/DCR.0b013e3181db7738]
- 30 **Naimy N**, Lindam AT, Bakka A, Faerden AE, Wiik P, Carlsen E, Nesheim BI. Biofeedback vs. electrostimulation in the treatment of postdelivery anal incontinence: a randomized, clinical trial. *Dis Colon Rectum* 2007; **50**: 2040-2046 [PMID: 17914654 DOI: 10.1007/s10350-007-9075-5]
- 31 **Jodorkovsky D**, Dunbar KB, Gearhart SL, Stein EM, Clarke JO. Biofeedback therapy for defecatory dysfunction: "real life" experience. *J Clin Gastroenterol* 2013; **47**: 252-255 [PMID: 23328298 DOI: 10.1097/MCG.0b013e318266f43a]
- 32 **Byrne CM**, Solomon MJ, Young JM, Rex J, Merlino CL. Biofeedback for fecal incontinence: short-term outcomes of 513 consecutive patients and predictors of successful treatment. *Dis Colon Rectum* 2007; **50**: 417-427 [PMID: 17476558 DOI: 10.1007/s10350-006-0846-1]
- 33 **Keck JO**, Staniunas RJ, Collier JA, Barrett RC, Oster ME, Schoetz DJ, Roberts PL, Murray JJ, Veidenheimer MC. Biofeedback training is useful in fecal incontinence but disappointing in constipation. *Dis Colon Rectum* 1994; **37**: 1271-1276 [PMID: 7995157 DOI: 10.1007/BF02257795]
- 34 **Terra MP**, Dobben AC, Berghmans B, Deutekom M, Baeten CG, Janssen LW, Boeckxstaens GE, Engel AF, Felt-Bersma RJ, Slors JF, Gerhards MF, Bijnen AB, Everhardt E, Schouten WR, Bossuyt PM, Stoker J. Electrical stimulation and pelvic floor muscle training with biofeedback in patients with fecal incontinence: a cohort study of 281 patients. *Dis Colon Rectum* 2006; **49**: 1149-1159 [PMID: 16773492 DOI: 10.1007/s10350-006-0569-3]
- 35 **Norton C**, Chelvanayagam S, Wilson-Barnett J, Redfern S, Kamm MA. Randomized controlled trial of biofeedback for fecal incontinence. *Gastroenterology* 2003; **125**: 1320-1329 [PMID: 14598248]
- 36 **Goetz LH**, Lowry AC. Overlapping sphincteroplasty: is it the standard of care? *Clin Colon Rectal Surg* 2005; **18**: 22-31 [PMID: 20011336 DOI: 10.1055/s-2005-864072]
- 37 **Oliveira L**, Pfeifer J, Wexner SD. Physiological and clinical outcome of anterior sphincteroplasty. *Br J Surg* 1996; **83**: 502-505 [PMID: 8665241 DOI: 10.1002/bjs.1800830421]
- 38 **Johnson E**, Carlsen E, Steen TB, Backer Hjorthaug JO, Eriksen MT, Johannessen HO. Short- and long-term results of secondary anterior sphincteroplasty in 33 patients with obstetric injury. *Acta Obstet Gynecol Scand* 2010; **89**: 1466-1472 [PMID: 20955101 DOI: 10.3109/00016349.2010.519019]
- 39 **Bravo Gutierrez A**, Madoff RD, Lowry AC, Parker SC, Buie WD, Baxter NN. Long-term results of anterior sphincteroplasty. *Dis Colon Rectum* 2004; **47**: 727-731; discussion 731-732 [PMID: 15037931 DOI: 10.1007/s10350-003-0114-6]
- 40 **Halverson AL**, Hull TL. Long-term outcome of overlapping anal sphincter repair. *Dis Colon Rectum* 2002; **45**: 345-348 [PMID: 12068192 DOI: 10.1007/s10350-004-6180-6]
- 41 **Malouf AJ**, Norton CS, Engel AF, Nicholls RJ, Kamm MA. Long-term results of overlapping anterior anal-sphincter repair for obstetric trauma. *Lancet* 2000; **355**: 260-265 [PMID: 10675072]
- 42 **Oom DM**, Steensma AB, Zimmerman DD, Schouten WR. Anterior sphincteroplasty for fecal incontinence: is the outcome compromised in patients with associated pelvic floor injury? *Dis Colon Rectum* 2010; **53**: 150-155 [PMID: 20087089 DOI: 10.1007/DCR.0b013e3181bb059f]
- 43 **Oberwalder M**, Dinnewitzer A, Baig MK, Noguerras JJ, Weiss EG, Efron J, Vernava AM, Wexner SD. Do internal anal sphincter defects decrease the success rate of anal sphincter repair? *Tech Coloproctol* 2006; **10**: 94-97; discussion 97 [PMID: 16773293 DOI: 10.1007/s10151-006-0259-0]
- 44 **Briel JW**, de Boer LM, Hop WC, Schouten WR. Clinical outcome of anterior overlapping external anal sphincter repair with internal anal sphincter imbrication. *Dis Colon Rectum* 1998; **41**: 209-214 [PMID: 9556246 DOI: 10.1007/BF02238250]
- 45 **Soerensen MM**, Bek KM, Buntzen S, Højberg KE, Laurberg S. Long-term outcome of delayed primary or early secondary reconstruction of the anal sphincter after obstetrical injury. *Dis Colon Rectum* 2008; **51**: 312-317 [PMID: 18219533 DOI: 10.1007/s10350-007-9084-4]
- 46 **Tan EK**, Jacovides M, Khullar V, Teoh TG, Fernando RJ, Tekkis PP. A cost-effectiveness analysis of delayed sphincteroplasty for anal sphincter injury. *Colorectal Dis* 2008; **10**: 653-662 [PMID: 18355373 DOI: 10.1111/j.1463-1318.2008.01507.x]
- 47 **Matsuoka H**, Mavrantonis C, Wexner SD, Oliveira L, Gilliland R, Pikarsky A. Postanal repair for fecal incontinence—is it worthwhile? *Dis Colon Rectum* 2000; **43**: 1561-1567 [PMID: 11089593 DOI: 10.1007/BF02236739]
- 48 **Orrom WJ**, Miller R, Cornes H, Duthie G, Mortensen NJ, Bartolo DC. Comparison of anterior sphincteroplasty and postanal repair in the treatment of idiopathic fecal incontinence. *Dis Colon Rectum* 1991; **34**: 305-310 [PMID: 2007347 DOI: 10.1007/BF02050589]
- 49 **Barisic GI**, Krivokapic ZV, Markovic VA, Popovic MA. Outcome of overlapping anal sphincter repair after 3 months and after a mean of 80 months. *Int J Colorectal Dis* 2006; **21**: 52-56 [PMID: 15830204 DOI: 10.1007/s00384-004-0737-7]
- 50 **Oom DM**, Gosselink MP, Schouten WR. Anterior sphincteroplasty for fecal incontinence: a single center experience in the era of sacral neuromodulation. *Dis Colon Rectum* 2009; **52**: 1681-1687 [PMID: 19966598 DOI: 10.1007/DCR.0b013e3181b13862]
- 51 **Mevik K**, Norderval S, Kileng H, Johansen M, Vonen B. Long-term results after anterior sphincteroplasty for anal incontinence. *Scand J Surg* 2009; **98**: 234-238 [PMID: 20218421]
- 52 **Karoui S**, Leroi AM, Koning E, Menard JF, Michot F, Denis P. Results of sphincteroplasty in 86 patients with anal incontinence. *Dis Colon Rectum* 2000; **43**: 813-820 [PMID: 10859083 DOI: 10.1007/BF02238020]
- 53 **Zutshi M**, Tracey TH, Bast J, Halverson A, Na J. Ten-year outcome after anal sphincter repair for fecal incontinence. *Dis Colon Rectum* 2009; **52**: 1089-1094 [PMID: 19581851 DOI: 10.1007/DCR.0b013e3181a0a79c]
- 54 **Glasgow SC**, Lowry AC. Long-term outcomes of anal sphincter repair for fecal incontinence: a systematic review. *Dis Colon Rectum* 2012; **55**: 482-490 [PMID: 22426274 DOI: 10.1097/DCR.0b013e3182468c22]
- 55 **Maslekar S**, Gardiner AB, Duthie GS. Anterior anal sphincter repair for fecal incontinence: Good longterm results are possible. *J Am Coll Surg* 2007; **204**: 40-46 [PMID: 17189111 DOI: 10.1016/j.jamcollsurg.2006.10.008]
- 56 **Lehto K**, Hyöty M, Collin P, Huhtala H, Aitola P. Seven-year follow-up after anterior sphincter reconstruction for faecal incontinence. *Int J Colorectal Dis* 2013; **28**: 653-658 [PMID: 23440365 DOI: 10.1007/s00384-013-1663-3]
- 57 **El-Gazzaz G**, Zutshi M, Hannaway C, Gurland B, Hull T. Overlapping sphincter repair: does age matter? *Dis Colon Rectum* 2012; **55**: 256-261 [PMID: 22469791 DOI: 10.1097/DCR.0b013e31823deb85]
- 58 **Altomare DF**, De Fazio M, Giuliani RT, Catalano G, Cuccia F. Sphincteroplasty for fecal incontinence in the era of sacral nerve modulation. *World J Gastroenterol* 2010; **16**: 5267-5271 [PMID: 21072888 DOI: 10.3748/wjg.v16.i42.5267]

- 59 **Gallas S**, Michot F, Faucheron JL, Meurette G, Lehur PA, Barth X, Damon H, Mion F, Rullier E, Zerbib F, Sielezneff I, Ouaïssi M, Orsoni P, Desfourneaux V, Siproudhis L, Mathonnet M, Menard JF, Leroi AM. Predictive factors for successful sacral nerve stimulation in the treatment of faecal incontinence: results of trial stimulation in 200 patients. *Colorectal Dis* 2011; **13**: 689-696 [PMID: 20236144 DOI: 10.1111/j.1463-1318.2010.02260.x]
- 60 **Ratto C**, Litta F, Parello A, Donisi L, De Simone V, Zaccone G. Sacral nerve stimulation in faecal incontinence associated with an anal sphincter lesion: a systematic review. *Colorectal Dis* 2012; **14**: e297-e304 [PMID: 22356165 DOI: 10.1111/j.1463-1318.2012.03003.x]
- 61 **Iachetta RP**, Cola A, Villani RD. Sacral nerve stimulation in the treatment of fecal incontinence - the experience of a pelvic floor center : short term results. *J Interv Gastroenterol* 2012; **2**: 189-192 [PMID: 23687607]
- 62 **Ratto C**, Litta F, Parello A, Donisi L, Doglietto GB. Sacral nerve stimulation is a valid approach in fecal incontinence due to sphincter lesions when compared to sphincter repair. *Dis Colon Rectum* 2010; **53**: 264-272 [PMID: 20173471 DOI: 10.1007/DCR.0b013e3181c7642c]
- 63 **Boyle DJ**, Knowles CH, Lunniss PJ, Scott SM, Williams NS, Gill KA. Efficacy of sacral nerve stimulation for fecal incontinence in patients with anal sphincter defects. *Dis Colon Rectum* 2009; **52**: 1234-1239 [PMID: 19571698 DOI: 10.1007/DCR.0b013e31819f7400]
- 64 **Brouwer R**, Duthie G. Sacral nerve neuromodulation is effective treatment for fecal incontinence in the presence of a sphincter defect, pudendal neuropathy, or previous sphincter repair. *Dis Colon Rectum* 2010; **53**: 273-278 [PMID: 20173472 DOI: 10.1007/DCR.0b013e3181cceb22]
- 65 **Gourcerol G**, Vitton V, Leroi AM, Michot F, Abysique A, Bouvier M. How sacral nerve stimulation works in patients with faecal incontinence. *Colorectal Dis* 2011; **13**: e203-e211 [PMID: 21689312 DOI: 10.1111/j.1463-1318.2011.02623.x]
- 66 **Brill SA**, Margolin DA. Sacral nerve stimulation for the treatment of fecal incontinence. *Clin Colon Rectal Surg* 2005; **18**: 38-41 [PMID: 20011338 DOI: 10.1055/s-2005-864079]
- 67 **Otto SD**, Burmeister S, Buhr HJ, Kroesen A. Sacral nerve stimulation induces changes in the pelvic floor and rectum that improve continence and quality of life. *J Gastrointest Surg* 2010; **14**: 636-644 [PMID: 20058096 DOI: 10.1007/s11605-009-1122-1]
- 68 **Patton V**, Wiklendt L, Arkwright JW, Lubowski DZ, Dinning PG. The effect of sacral nerve stimulation on distal colonic motility in patients with faecal incontinence. *Br J Surg* 2013; **100**: 959-968 [PMID: 23536312 DOI: 10.1002/bjs.9114]
- 69 **Griffin KM**, Pickering M, O'Herlihy C, O'Connell PR, Jones JF. Sacral nerve stimulation increases activation of the primary somatosensory cortex by anal canal stimulation in an experimental model. *Br J Surg* 2011; **98**: 1160-1169 [PMID: 21590761 DOI: 10.1002/bjs.7536]
- 70 **Vallet C**, Parc Y, Lupinacci R, Shields C, Parc R, Tiret E. Sacral nerve stimulation for faecal incontinence: response rate, satisfaction and the value of preoperative investigation in patient selection. *Colorectal Dis* 2010; **12**: 247-253 [PMID: 19508523 DOI: 10.1111/j.1463-1318.2009.01899.x]
- 71 **Norderval S**, Behrenbruch C, Brouwer R, Keck JO. Efficacy of cyclic sacral nerve stimulation for faecal incontinence. *Tech Coloproctol* 2013; **17**: 511-516 [PMID: 23525966]
- 72 **Duelund-Jakobsen J**, Buntzen S, Lundby L, Laurberg S. Sacral nerve stimulation at subsensory threshold does not compromise treatment efficacy: results from a randomized, blinded crossover study. *Ann Surg* 2013; **257**: 219-223 [PMID: 23001079 DOI: 10.1097/SLA.0b013e318269d493]
- 73 **Leroi AM**, Parc Y, Lehur PA, Mion F, Barth X, Rullier E, Bresler L, Portier G, Michot F. Efficacy of sacral nerve stimulation for fecal incontinence: results of a multicenter double-blind crossover study. *Ann Surg* 2005; **242**: 662-669 [PMID: 16244539 DOI: 10.1097/01.sla.0000186281.09475.db]
- 74 **Tan E**, Ngo NT, Darzi A, Shenouda M, Tekkis PP. Meta-analysis: sacral nerve stimulation versus conservative therapy in the treatment of faecal incontinence. *Int J Colorectal Dis* 2011; **26**: 275-294 [PMID: 21279370 DOI: 10.1007/s00384-010-1119-y]
- 75 **Hull T**, Giese C, Wexner SD, Mellgren A, Devroede G, Madoff RD, Stromberg K, Collier JA. Long-term durability of sacral nerve stimulation therapy for chronic fecal incontinence. *Dis Colon Rectum* 2013; **56**: 234-245 [PMID: 23303153 DOI: 10.1097/DCR.0b013e318276b24c]
- 76 **Hollingshead JR**, Dudding TC, Vaizey CJ. Sacral nerve stimulation for faecal incontinence: results from a single centre over a 10-year period. *Colorectal Dis* 2011; **13**: 1030-1034 [PMID: 20718837 DOI: 10.1111/j.1463-1318.2010.02383.x]
- 77 **George AT**, Kalmar K, Panarese A, Dudding TC, Nicholls RJ, Vaizey CJ. Long-term outcomes of sacral nerve stimulation for fecal incontinence. *Dis Colon Rectum* 2012; **55**: 302-306 [PMID: 22469797 DOI: 10.1097/DCR.0b013e3182401ecd]
- 78 **Lim JT**, Hastie IA, Hiscock RJ, Shedda SM. Sacral nerve stimulation for fecal incontinence: long-term outcomes. *Dis Colon Rectum* 2011; **54**: 969-974 [PMID: 21730785 DOI: 10.1097/DCR.0b013e31821e57c2]
- 79 **Wexner SD**, Collier JA, Devroede G, Hull T, McCallum R, Chan M, Ayscue JM, Shobeiri AS, Margolin D, England M, Kaufman H, Snape WJ, Mutlu E, Chua H, Pettit P, Nagle D, Madoff RD, Lerew DR, Mellgren A. Sacral nerve stimulation for fecal incontinence: results of a 120-patient prospective multicenter study. *Ann Surg* 2010; **251**: 441-449 [PMID: 20160636 DOI: 10.1097/SLA.0b013e3181cf8ed0]
- 80 **Jadav AM**, Wadhawan H, Jones GL, Wheldon LW, Radley SC, Brown SR. Does sacral nerve stimulation improve global pelvic function in women? *Colorectal Dis* 2013; **15**: 848-857 [PMID: 23451900 DOI: 10.1111/codi.12181]
- 81 **Mellgren A**, Wexner SD, Collier JA, Devroede G, Lerew DR, Madoff RD, Hull T. Long-term efficacy and safety of sacral nerve stimulation for fecal incontinence. *Dis Colon Rectum* 2011; **54**: 1065-1075 [PMID: 21825885 DOI: 10.1097/DCR.0b013e31822155e9]
- 82 **Damon H**, Barth X, Roman S, Mion F. Sacral nerve stimulation for fecal incontinence improves symptoms, quality of life and patients' satisfaction: results of a monocentric series of 119 patients. *Int J Colorectal Dis* 2013; **28**: 227-233 [PMID: 22885883 DOI: 10.1007/s00384-012-1558-8]
- 83 **Devroede G**, Giese C, Wexner SD, Mellgren A, Collier JA, Madoff RD, Hull T, Stromberg K, Iyer S. Quality of life is markedly improved in patients with fecal incontinence after sacral nerve stimulation. *Female Pelvic Med Reconstr Surg* 2012; **18**: 103-112 [PMID: 22453321 DOI: 10.1097/SPV.0b013e3182486e60]
- 84 **Duelund-Jakobsen J**, van Wunnik B, Buntzen S, Lundby L, Baeten C, Laurberg S. Functional results and patient satisfaction with sacral nerve stimulation for idiopathic faecal incontinence. *Colorectal Dis* 2012; **14**: 753-759 [PMID: 21883814 DOI: 10.1111/j.1463-1318.2011.02800.x]
- 85 **Maeda Y**, Matzel K, Lundby L, Buntzen S, Laurberg S. Post-operative issues of sacral nerve stimulation for fecal incontinence and constipation: a systematic literature review and treatment guideline. *Dis Colon Rectum* 2011; **54**: 1443-1460 [PMID: 21979192 DOI: 10.1097/DCR.0b013e318227f65d]
- 86 **Leroi AM**, Lenne X, Dervaux B, Chartier-Kastler E, Mauroy B, Normand LL, Grise P, Faucheron JL, Parc Y, Lehur PA, Mion F, Damon H, Barth X, Leriche A, Saussine C, Guy L, Haab F, Bresler L, Sarramon JP, Bensadoun H, Rullier E, Slim K, Sielezneff I, Mourey E, Ballanger P, Michot F. Outcome and cost analysis of sacral nerve modulation for treating urinary and/or fecal incontinence. *Ann Surg* 2011; **253**: 720-732 [PMID: 21475012 DOI: 10.1097/SLA.0b013e318210f1f4]
- 87 **Indinnimeo M**, Ratto C, Moschella CM, Fiore A, Brosa M, Giardina S. Sacral neuromodulation for the treatment of fecal incontinence: analysis of cost-effectiveness. *Dis Colon*

- Rectum* 2010; **53**: 1661-1669 [PMID: 21178862 DOI: 10.1007/DCR.0b013e3181f46309]
- 88 **Wexner SD**, Hull T, Edden Y, Collier JA, Devroede G, McCallum R, Chan M, Ayscue JM, Shobeiri AS, Margolin D, England M, Kaufman H, Snape WJ, Mutlu E, Chua H, Pettit P, Nagle D, Madoff RD, Lerew DR, Mellgren A. Infection rates in a large investigational trial of sacral nerve stimulation for fecal incontinence. *J Gastrointest Surg* 2010; **14**: 1081-1089 [PMID: 20354809 DOI: 10.1007/s11605-010-1177-z]
- 89 **Gregorczyk SG**. The current status of the Acticon Neosphincter. *Clin Colon Rectal Surg* 2005; **18**: 32-37 [PMID: 20011337 DOI: 10.1055/s-2005-864078]
- 90 **Wexner SD**, Jin HY, Weiss EG, Nogueras JJ, Li VK. Factors associated with failure of the artificial bowel sphincter: a study of over 50 cases from Cleveland Clinic Florida. *Dis Colon Rectum* 2009; **52**: 1550-1557 [PMID: 19690481 DOI: 10.1007/DCR.0b013e3181af62f8]
- 91 **Ortiz H**, Armendariz P, DeMiguel M, Solana A, Alós R, Roig JV. Prospective study of artificial anal sphincter and dynamic graciloplasty for severe anal incontinence. *Int J Colorectal Dis* 2003; **18**: 349-354 [PMID: 12774251 DOI: 10.1007/s00384-002-0472-x]
- 92 **Tan EK**, Vaizey C, Cornish J, Darzi A, Tekkis PP. Surgical strategies for faecal incontinence--a decision analysis between dynamic graciloplasty, artificial bowel sphincter and end stoma. *Colorectal Dis* 2008; **10**: 577-586 [PMID: 18005188 DOI: 10.1111/j.1463-1318.2007.01418.x]
- 93 **Lehur PA**, Roig JV, Duinslaeger M. Artificial anal sphincter: prospective clinical and manometric evaluation. *Dis Colon Rectum* 2000; **43**: 1100-1106 [PMID: 10950008]
- 94 **Altomare DF**, Dodi G, La Torre F, Romano G, Melega E, Rinaldi M. Multicentre retrospective analysis of the outcome of artificial anal sphincter implantation for severe faecal incontinence. *Br J Surg* 2001; **88**: 1481-1486 [PMID: 11683745 DOI: 10.1046/j.0007-1323.2001.01895.x]
- 95 **Devesa JM**, Rey A, Hervas PL, Halawa KS, Larrañaga I, Svidler L, Abaira V, Muriel A. Artificial anal sphincter: complications and functional results of a large personal series. *Dis Colon Rectum* 2002; **45**: 1154-1163 [PMID: 12352229 DOI: 10.1007/s10350-004-6382-y]
- 96 **Parker SC**, Spencer MP, Madoff RD, Jensen LL, Wong WD, Rothenberger DA. Artificial bowel sphincter: long-term experience at a single institution. *Dis Colon Rectum* 2003; **46**: 722-729 [PMID: 12794572 DOI: 10.1007/s10350-004-6647-5]
- 97 **Wong WD**, Congliosi SM, Spencer MP, Corman ML, Tan P, Opelka FG, Burnstein M, Nogueras JJ, Bailey HR, Devesa JM, Fry RD, Cagir B, Birnbaum E, Fleshman JW, Lawrence MA, Buie WD, Heine J, Edelstein PS, Gregorczyk S, Lehur PA, Michot F, Phang PT, Schoetz DJ, Potenti F, Tsai JY. The safety and efficacy of the artificial bowel sphincter for fecal incontinence: results from a multicenter cohort study. *Dis Colon Rectum* 2002; **45**: 1139-1153 [PMID: 12352228 DOI: 10.1007/s10350-004-6381-z]
- 98 **O'Brien PE**, Dixon JB, Skinner S, Laurie C, Khera A, Fonda D. A prospective, randomized, controlled clinical trial of placement of the artificial bowel sphincter (Acticon Neosphincter) for the control of fecal incontinence. *Dis Colon Rectum* 2004; **47**: 1852-1860 [PMID: 15622577 DOI: 10.1007/s10350-004-0717-6]
- 99 **Ruiz Carmona MD**, Alós Company R, Roig Vila JV, Solana Bueno A, Pla Martí V. Long-term results of artificial bowel sphincter for the treatment of severe faecal incontinence. Are they what we hoped for? *Colorectal Dis* 2009; **11**: 831-837 [PMID: 18662237 DOI: 10.1111/j.1463-1318.2008.01652.x]
- 100 **Wong MT**, Meurette G, Wyart V, Glemain P, Lehur PA. The artificial bowel sphincter: a single institution experience over a decade. *Ann Surg* 2011; **254**: 951-956 [PMID: 22107742 DOI: 10.1097/SLA.0b013e31823ac2bc]
- 101 **Lehur PA**, Zerbib F, Neunlist M, Glemain P, Bruley des Varannes S. Comparison of quality of life and anorectal function after artificial sphincter implantation. *Dis Colon Rectum* 2002; **45**: 508-513 [PMID: 12006933 DOI: 10.1007/s10350-004-6230-0]
- 102 **Mundy L**, Merlin TL, Maddern GJ, Hiller JE. Systematic review of safety and effectiveness of an artificial bowel sphincter for faecal incontinence. *Br J Surg* 2004; **91**: 665-672 [PMID: 15164433 DOI: 10.1002/bjs.4587]
- 103 **Melenhorst J**, Koch SM, van Gemert WG, Baeten CG. The artificial bowel sphincter for faecal incontinence: a single centre study. *Int J Colorectal Dis* 2008; **23**: 107-111 [PMID: 17929038 DOI: 10.1007/s00384-007-0357-0]
- 104 **Michot F**, Costaglioli B, Leroi AM, Denis P. Artificial anal sphincter in severe fecal incontinence: outcome of prospective experience with 37 patients in one institution. *Ann Surg* 2003; **237**: 52-56 [PMID: 12496530]
- 105 **Gallas S**, Leroi AM, Bridoux V, Lefebure B, Tuech JJ, Michot F. Constipation in 44 patients implanted with an artificial bowel sphincter. *Int J Colorectal Dis* 2009; **24**: 969-974 [PMID: 19283389 DOI: 10.1007/s00384-009-0693-3]
- 106 **Cera SM**, Wexner SD. Muscle transposition: does it still have a role? *Clin Colon Rectal Surg* 2005; **18**: 46-54 [PMID: 20011340 DOI: 10.1055/s-2005-864081]
- 107 **Hassan MZ**, Rathnayaka MM, Deen KI. Modified dynamic gracilis neosphincter for fecal incontinence: an analysis of functional outcome at a single institution. *World J Surg* 2010; **34**: 1641-1647 [PMID: 20180122 DOI: 10.1007/s00268-010-0489-1]
- 108 **Devesa JM**, Madrid JM, Gallego BR, Vicente E, Nuño J, Enriquez JM. Bilateral gluteoplasty for fecal incontinence. *Dis Colon Rectum* 1997; **40**: 883-888 [PMID: 9269802 DOI: 10.1007/BF02051193]
- 109 **Madoff RD**, Rosen HR, Baeten CG, LaFontaine LJ, Cavina E, Devesa M, Rouanet P, Christiansen J, Faucheron JL, Isbister W, Köhler L, Guelinckx PJ, Pählman L. Safety and efficacy of dynamic muscle plasty for anal incontinence: lessons from a prospective, multicenter trial. *Gastroenterology* 1999; **116**: 549-556 [PMID: 10029613 DOI: 10.1016/S0016-5085(99)70176-9]
- 110 **Wexner SD**, Baeten C, Bailey R, Bakka A, Belin B, Belliveau P, Berg E, Buie WD, Burnstein M, Christiansen J, Collier J, Galandiuk S, Lange J, Madoff R, Matzel KE, Pählman L, Parc R, Reilly J, Seccia M, Thorson AG, Vernava AM. Long-term efficacy of dynamic graciloplasty for fecal incontinence. *Dis Colon Rectum* 2002; **45**: 809-818 [PMID: 12072635 DOI: 10.1007/s10350-004-6302-1]
- 111 **Rongen MJ**, Uludag O, El Naggar K, Geerdes BP, Konsten J, Baeten CG. Long-term follow-up of dynamic graciloplasty for fecal incontinence. *Dis Colon Rectum* 2003; **46**: 716-721 [PMID: 12794571 DOI: 10.1007/s10350-004-6645-7]
- 112 **Bresler L**, Reibel N, Brunaud L, Sielezneff I, Rouanet P, Rullier E, Slim K. [Dynamic graciloplasty in the treatment of severe fecal incontinence. French multicentric retrospective study]. *Ann Chir* 2002; **127**: 520-526 [PMID: 12404846]
- 113 **Eccersley AJ**, Lunniss PJ, Williams NS. Unstimulated graciloplasty in traumatic faecal incontinence. *Br J Surg* 1999; **86**: 1071-1072 [PMID: 10460647 DOI: 10.1046/j.1365-2168.1999.01159.x]
- 114 **Kumar D**, Hutchinson R, Grant E. Bilateral gracilis neosphincter construction for treatment of faecal incontinence. *Br J Surg* 1995; **82**: 1645-1647 [PMID: 8548229 DOI: 10.1002/bjs.1800821219]
- 115 **Matzel KE**, Madoff RD, LaFontaine LJ, Baeten CG, Buie WD, Christiansen J, Wexner S. Complications of dynamic graciloplasty: incidence, management, and impact on outcome. *Dis Colon Rectum* 2001; **44**: 1427-1435 [PMID: 11598470 DOI: 10.1007/BF02234593]
- 116 **Rosen HR**, Urbarz C, Novi G, Zöch G, Schiessel R. Long-term results of modified graciloplasty for sphincter replacement after rectal excision. *Colorectal Dis* 2002; **4**: 266-269 [PMID: 12780598 DOI: 10.1046/j.1463-1318.2002.00331.x]
- 117 **Thornton MJ**, Kennedy ML, Lubowski DZ, King DW. Long-term follow-up of dynamic graciloplasty for faecal incontinence. *Colorectal Dis* 2004; **6**: 470-476 [PMID: 15521938 DOI:

- 10.1111/j.1463-1318.2004.00714.x]
- 118 **da Silva GM**, Jorge JM, Belin B, Noguerras JJ, Weiss EG, Vernava AM, Habr-Gama A, Wexner SD. New surgical options for fecal incontinence in patients with imperforate anus. *Dis Colon Rectum* 2004; **47**: 204-209 [PMID: 15043291 DOI: 10.1007/s10350-003-0039-0]
- 119 **Malone PS**, Ransley PG, Kiely EM. Preliminary report: the antegrade continence enema. *Lancet* 1990; **336**: 1217-1218 [PMID: 1978072 DOI: 10.1016/0140-6736(90)92834-5]
- 120 **Teichman JM**, Harris JM, Currie DM, Barber DB. Malone antegrade continence enema for adults with neurogenic bowel disease. *J Urol* 1998; **160**: 1278-1281 [PMID: 9751335 DOI: 10.1016/S0022-5347(01)62515-1]
- 121 **Gerharz EW**, Vik V, Webb G, Leaver R, Shah PJ, Woodhouse CR. The value of the MACE (Malone antegrade colonic enema) procedure in adult patients. *J Am Coll Surg* 1997; **185**: 544-547 [PMID: 9404877]
- 122 **Tackett LD**, Minevich E, Benedict JF, Wacksman J, Sheldon CA. Appendiceal versus ileal segment for antegrade continence enema. *J Urol* 2002; **167**: 683-686 [PMID: 11792954 DOI: 10.1016/S0022-5347(01)69125-0]
- 123 **Karpman E**, Das S, Kurzrock EA. Laparoscopic antegrade continence enema (Malone) procedure: description and illustration of technique. *J Endourol* 2002; **16**: 325-328; discussion 328 [PMID: 12227902 DOI: 10.1089/089277902760261310]
- 124 **Ellison JS**, Haraway AN, Park JM. The distal left Malone antegrade continence enema--is it better? *J Urol* 2013; **190**: 1529-1533 [PMID: 23376708 DOI: 10.1016/j.juro.2013.01.092]
- 125 **Bani-Hani AH**, Cain MP, King S, Rink RC. Tap water irrigation and additives to optimize success with the Malone antegrade continence enema: the Indiana University algorithm. *J Urol* 2008; **180**: 1757-1760; discussion 1760 [PMID: 18721951 DOI: 10.1016/j.juro.2008.04.074]
- 126 **Poirier M**, Abcarian H, Nelson R. Malone antegrade continence enema: an alternative to resection in severe defecation disorders. *Dis Colon Rectum* 2007; **50**: 22-28 [PMID: 17115341 DOI: 10.1007/s10350-006-0732-x]
- 127 **Lefèvre JH**, Parc Y, Giraudo G, Bell S, Parc R, Turet E. Outcome of antegrade continence enema procedures for faecal incontinence in adults. *Br J Surg* 2006; **93**: 1265-1269 [PMID: 16952210 DOI: 10.1002/bjs.5383]
- 128 **Teichman JM**, Zabihi N, Kraus SR, Harris JM, Barber DB. Long-term results for Malone antegrade continence enema for adults with neurogenic bowel disease. *Urology* 2003; **61**: 502-506 [PMID: 12639632 DOI: 10.1016/S0090-4295(02)02282-3]
- 129 **Bani-Hani AH**, Cain MP, Kaefer M, Meldrum KK, King S, Johnson CS, Rink RC. The Malone antegrade continence enema: single institutional review. *J Urol* 2008; **180**: 1106-1110 [PMID: 18639902 DOI: 10.1016/j.juro.2008.05.062]
- 130 **Herndon CD**, Rink RC, Cain MP, Lerner M, Kaefer M, Yerkes E, Casale AJ. In situ Malone antegrade continence enema in 127 patients: a 6-year experience. *J Urol* 2004; **172**: 1689-1691 [PMID: 15371791 DOI: 10.1016/S0022-5347(05)61339-0]
- 131 **Yerkes EB**, Cain MP, King S, Brei T, Kaefer M, Casale AJ, Rink RC. The Malone antegrade continence enema procedure: quality of life and family perspective. *J Urol* 2003; **169**: 320-323 [PMID: 12478181 DOI: 10.1016/S0022-5347(05)64116-X]
- 132 **Soylet Y**, Yesildag E, Besik C, Emir H. Antegrade continence enema--an analysis of 20 children with faecal incontinence. *Eur J Pediatr Surg* 2006; **16**: 251-254 [PMID: 16981089 DOI: 10.1055/s-2006-924337]
- 133 **Nanigian DK**, Kurzrock EA. Intermediate-term outcome of the simplified laparoscopic antegrade continence enema procedure: less is better. *J Urol* 2008; **179**: 299-303 [PMID: 18006019 DOI: 10.1016/j.juro.2007.08.161]
- 134 **Ok JH**, Kurzrock EA. Objective measurement of quality of life changes after ACE Malone using the FICQOL survey. *J Pediatr Urol* 2011; **7**: 389-393 [PMID: 21527223 DOI: 10.1016/j.jpuro.2011.02.012]
- 135 **Tiryaki S**, Ergun O, Celik A, Ulman I, Avanoğlu A. Success of Malone's antegrade continence enema (MACE) from the patients' perspective. *Eur J Pediatr Surg* 2010; **20**: 405-407 [PMID: 20954103 DOI: 10.1055/s-0030-1265156]
- 136 **Mattix KD**, Novotny NM, Shelley AA, Rescorla FJ. Malone antegrade continence enema (MACE) for fecal incontinence in imperforate anus improves quality of life. *Pediatr Surg Int* 2007; **23**: 1175-1177 [PMID: 17938937 DOI: 10.1007/s00383-007-2026-3]
- 137 **Van Savage JG**, Yohannes P. Laparoscopic antegrade continence enema in situ appendix procedure for refractory constipation and overflow fecal incontinence in children with spina bifida. *J Urol* 2000; **164**: 1084-1087 [PMID: 10958747 DOI: 10.1016/S0022-5347(05)67257-6]
- 138 **Hoekstra LT**, Kuijper CF, Bakx R, Heij HA, Aronson DC, Benninga MA. The Malone antegrade continence enema procedure: the Amsterdam experience. *J Pediatr Surg* 2011; **46**: 1603-1608 [PMID: 21843730 DOI: 10.1016/j.jpedsurg.2011.04.050]
- 139 **Colquhoun P**, Kaiser R, Efron J, Weiss EG, Noguerras JJ, Vernava AM, Wexner SD. Is the quality of life better in patients with colostomy than patients with fecal incontinence? *World J Surg* 2006; **30**: 1925-1928 [PMID: 16957817 DOI: 10.1007/s00268-006-0531-5]
- 140 **Norton C**, Burch J, Kamm MA. Patients' views of a colostomy for fecal incontinence. *Dis Colon Rectum* 2005; **48**: 1062-1069 [PMID: 15868244 DOI: 10.1007/s10350-004-0868-5]
- 141 **Parisien CJ**, Corman ML. The Secca procedure for the treatment of fecal incontinence: definitive therapy or short-term solution. *Clin Colon Rectal Surg* 2005; **18**: 42-45 [PMID: 20011339 DOI: 10.1055/s-2005-864080]
- 142 **Takahashi-Monroy T**, Morales M, Garcia-Osogobio S, Valdovinos MA, Belmonte C, Barreto C, Zarate X, Bada O, Velasco L. SECCA procedure for the treatment of fecal incontinence: results of five-year follow-up. *Dis Colon Rectum* 2008; **51**: 355-359 [PMID: 18204954 DOI: 10.1007/s10350-007-9169-0]
- 143 **Efron JE**, Corman ML, Fleshman J, Barnett J, Nagle D, Birnbaum E, Weiss EG, Noguerras JJ, Sligh S, Rabine J, Wexner SD. Safety and effectiveness of temperature-controlled radiofrequency energy delivery to the anal canal (Secca procedure) for the treatment of fecal incontinence. *Dis Colon Rectum* 2003; **46**: 1606-116; discussion 1606-116; [PMID: 14668584 DOI: 10.1007/BF02660763]
- 144 **Lefebvre B**, Tuech JJ, Bridoux V, Gallas S, Leroi AM, Denis P, Michot F. Temperature-controlled radio frequency energy delivery (Secca procedure) for the treatment of fecal incontinence: results of a prospective study. *Int J Colorectal Dis* 2008; **23**: 993-997 [PMID: 18594840 DOI: 10.1007/s00384-008-0514-0]
- 145 **Ruiz D**, Pinto RA, Hull TL, Efron JE, Wexner SD. Does the radiofrequency procedure for fecal incontinence improve quality of life and incontinence at 1-year follow-up? *Dis Colon Rectum* 2010; **53**: 1041-1046 [PMID: 20551757 DOI: 10.1007/DCR.0b013e3181deff8]
- 146 **Abbas MA**, Tam MS, Chun LJ. Radiofrequency treatment for fecal incontinence: is it effective long-term? *Dis Colon Rectum* 2012; **55**: 605-610 [PMID: 22513440 DOI: 10.1097/DCR.0b013e3182415406]
- 147 **Felt-Bersma RJ**, Szojda MM, Mulder CJ. Temperature-controlled radiofrequency energy (SECCA) to the anal canal for the treatment of faecal incontinence offers moderate improvement. *Eur J Gastroenterol Hepatol* 2007; **19**: 575-580 [PMID: 17556904 DOI: 10.1097/MEG.0b013e32811ec010]
- 148 **Kim DW**, Yoon HM, Park JS, Kim YH, Kang SB. Radiofrequency energy delivery to the anal canal: is it a promising new approach to the treatment of fecal incontinence? *Am J Surg* 2009; **197**: 14-18 [PMID: 18614149 DOI: 10.1016/j.amjsurg.2007.11.023]
- 149 **Watson NF**, Koshy A, Sagar PM. Anal bulking agents for faecal incontinence. *Colorectal Dis* 2012; **14** Suppl 3: 29-33 [PMID: 23136822 DOI: 10.1111/codi.12047]
- 150 **Dodi G**, Jongen J, de la Portilla F, Raval M, Altomare DF, Lehur PA. An Open-Label, Noncomparative, Multicenter

- Study to Evaluate Efficacy and Safety of NASHA/Dx Gel as a Bulking Agent for the Treatment of Fecal Incontinence. *Gastroenterol Res Pract* 2010; **2010**: 467136 [PMID: 21234379 DOI: 10.1155/2010/467136]
- 151 **Tjandra JJ**, Chan MK, Yeh HC. Injectable silicone biomaterial (PTQ) is more effective than carbon-coated beads (Durasphere) in treating passive faecal incontinence—a randomized trial. *Colorectal Dis* 2009; **11**: 382-389 [PMID: 18637935 DOI: 10.1111/j.1463-1318.2008.01634.x]
- 152 **Dehli T**, Stordahl A, Vatten LJ, Romundstad PR, Mevik K, Sahlin Y, Lindsetmo RO, Vonen B. Sphincter training or anal injections of dextranomer for treatment of anal incontinence: a randomized trial. *Scand J Gastroenterol* 2013; **48**: 302-310 [PMID: 23298304 DOI: 10.3109/00365521.2012.758770]
- 153 **Oliveira LC**, Neves Jorge JM, Yussuf S, Habr-Gama A, Kiss D, Ceconello I. Anal incontinence improvement after silicone injection may be related to restoration of sphincter asymmetry. *Surg Innov* 2009; **16**: 155-161 [PMID: 19546122 DOI: 10.1177/1553350609338374]
- 154 **Altomare DF**, La Torre F, Rinaldi M, Binda GA, Pescatori M. Carbon-coated microbeads anal injection in outpatient treatment of minor fecal incontinence. *Dis Colon Rectum* 2008; **51**: 432-435 [PMID: 18204878 DOI: 10.1007/s10350-007-9170-7]
- 155 **Danielson J**, Karlbom U, Wester T, Graf W. Efficacy and quality of life 2 years after treatment for faecal incontinence with injectable bulking agents. *Tech Coloproctol* 2013; **17**: 389-395 [PMID: 23224913 DOI: 10.1007/s10151-012-0949-8]
- 156 **La Torre F**, de la Portilla F. Long-term efficacy of dextranomer in stabilized hyaluronic acid (NASHA/Dx) for treatment of faecal incontinence. *Colorectal Dis* 2013; **15**: 569-574 [PMID: 23374680 DOI: 10.1111/codi.12155]
- 157 **Maeda Y**, Laurberg S, Norton C. Perianal injectable bulking agents as treatment for faecal incontinence in adults. *Cochrane Database Syst Rev* 2013; **2**: CD007959 [PMID: 23450581 DOI: 10.1002/14651858.CD007959.pub3]
- 158 **Tjandra JJ**, Lim JF, Hiscock R, Rajendra P. Injectable silicone biomaterial for fecal incontinence caused by internal anal sphincter dysfunction is effective. *Dis Colon Rectum* 2004; **47**: 2138-2146 [PMID: 15657666 DOI: 10.1007/s10350-004-0760-3]
- 159 **Schwandner O**, Brunner M, Dietl O. Quality of life and functional results of submucosal injection therapy using dextranomer hyaluronic acid for fecal incontinence. *Surg Innov* 2011; **18**: 130-135 [PMID: 21245071 DOI: 10.1177/1553350610392243]
- 160 **Danielson J**, Karlbom U, Sonesson AC, Wester T, Graf W. Submucosal injection of stabilized nonanimal hyaluronic acid with dextranomer: a new treatment option for fecal incontinence. *Dis Colon Rectum* 2009; **52**: 1101-1106 [PMID: 19581853 DOI: 10.1007/DCR.0b013e31819f5cbf]
- 161 **Graf W**, Mellgren A, Matzel KE, Hull T, Johansson C, Bernstein M. Efficacy of dextranomer in stabilised hyaluronic acid for treatment of faecal incontinence: a randomised, sham-controlled trial. *Lancet* 2011; **377**: 997-1003 [PMID: 21420555 DOI: 10.1016/S0140-6736(10)62297-0]

P- Reviewer: Maglinte DDT S- Editor: Zhai HH L- Editor: A
E- Editor: Ma S





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza,

315-321 Lockhart Road, Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045

